Food Allergy Action Plan

	Student's
re reaction)	Picture Here
the allergen was <i>likely</i> eat efinitely eaten, even if no s	
asthma	ELY pring (see box nal medications:* ne nchodilator) if halers/bronchodilators ded upon to treat a
2 parent 3. If symptoms	dent; alert rofessionals and progress (see EPINEPHRINE
	 the allergen was <i>likely</i> eat efinitely eaten, even if no s 1. INJECT EP IMMEDIATE 2. Call 911 3. Begin monito below) 4. Give addition -Antihistamin -Inhaler (bro asthma *Antihistamines & in are not to be depend severe reaction (ana EPINEPHRINE. 1. GIVE ANTIH 2. Stay with stur healthcare p parent 3. If symptoms above), USE 4. Begin monitor

Antihistamine (brand and dose):

Other (e.g., inhaler-bronchodilator if asthmatic):

Monitoring

Stay with student; alert healthcare professionals and parent. Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping student lying on back with legs raised. Treat student even if parents cannot be reached. See back/attached for auto-injection technique.

Parent/Guardian Signature

Date